

*Equivalent Characterizations of Lattice
Supersolvability and Their Extensions*

Peter McNamara

Joint work with
Hugh Thomas

Joint Mathematics Meetings
January 18th 2003

Slides and preprints available from
<http://www-math.mit.edu/~mcnamara/>


We say that a lattice L is *distributive* if

$$x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$$


and

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$$

for all elements x, y and z of L .



EXAMPLE The lattice of *order ideals* of a poset P .



An edge-labelling of a poset P is said to be an S_n *EL-labelling* if it satisfies the following 2 conditions:

1. Every interval $[x, y]$ of P has exactly one maximal chain with increasing labels
2. The labels along any maximal chain form a permutation of n .

Special case of *EL-labelling* (Björner):

2. The sequence of labels along this increasing maximal chain lexicographically precede the labels along any other maximal chain of $[x, y]$.


Who cares?

- EL-labelling \Rightarrow Shellable \Rightarrow Cohen-Macaulay
- Simple combinatorial interpretations of Möbius function, flag h-vector, etc.

What other classes of posets have S_n EL-labellings?

Definition(R. Stanley, 1972) A finite lattice L is said to be *supersolvable* if it contains a maximal chain \mathfrak{m} , called an *M -chain* of L , which together with any other chain of L generates a distributive sublattice.


EXAMPLES



- Distributive lattices
- The lattice of partitions of $\{1, 2, \dots, n\}$
- The lattice of non-crossing partitions of $\{1, 2, \dots, n\}$
- The lattice of subgroups of a supersolvable group

QUESTION (Stanley) Are there any other lattices that have S_n EL-labellings?

THEOREM (McN.) *A finite lattice has an S_n EL-labelling if and only if it is supersolvable.*




Connections with modularity...

Definition An element x of a lattice L is said to be *left-modular* if, for all $y \leq z$ in L , we have

$$(x \vee y) \wedge z = (x \wedge z) \vee y.$$

A chain of L is *left-modular* if each of its elements is left-modular.

Suppose L is a graded lattice.



L has a left-modular maximal chain

↙ Stanley

Liu ↖

THEOREM *Let L be graded lattice. TFAE:*


1. *L is supersolvable*
2. *L has an S_n EL-labelling*
3. *L has a left-modular maximal chain*
- 4.

How can we extend this?

- 3: L need not be graded
- 2: L need not be a lattice

Definition Let P be a (bounded) poset. An EL-labelling γ of P is said to be *interpolating* if, for any $y \triangleleft u \triangleleft z$, either

- (i) $\gamma(y, u) < \gamma(u, z)$ or
- (ii) the increasing chain from y to z , say $y = w_0 \triangleleft w_1 \triangleleft \cdots \triangleleft w_r = z$, has the properties that its labels are strictly increasing and that $\gamma(w_0, w_1) = \gamma(u, z)$ and $\gamma(w_{r-1}, w_r) = \gamma(y, u)$.




THEOREM (Thomas) *A lattice has an interpolating EL-labelling if and only if it has a left modular maximal chain.*

Generalizing to non-lattices:

P : a bounded poset with an S_n EL-labelling.

M : its increasing maximal chain.

Some “left modularity” property ?



When $x \in M$, $x \vee y$ and $x \wedge y$ are well-defined.

In a lattice: $(x \vee y) \wedge z \geq y$ whenever $z \geq y$.

When $x \in M$, $(x \vee y) \wedge_y z$ is well-defined for $y \leq z$. Similarly, $(x \wedge z) \vee^z y$ is well-defined.

We call x a *viable* element of P .

We call M a *viable* maximal chain.

THEOREM (McN.-Thomas) *A bounded poset has an interpolating EL-labelling if and only if it has a viable left modular maximal chain.*

	Graded	Not nec. graded
Lattice	1. Supersolvable 2. S_n EL-labelling 3. Left mod. max. chain	1. ? 2. Interp. EL-labelling 3. Left mod. max. chain
Not nec. Lattice	1. "Supersolvable" 2. S_n EL-labelling 3. Viable left mod. m.c.	1. ? 2. Interp. EL-labelling 3. Viable left mod. m.c.