

1052-05-58

**Bruce E. Sagan\*** ([sagan@math.msu.edu](mailto:sagan@math.msu.edu)), Department of Mathematics, East Lansing, MI 20912.  
*Probabilistic proofs of hook length formulas involving trees.*

Let  $T$  be a rooted tree on  $n$  vertices. We use  $T$  to stand for the vertex set of  $T$ . An *increasing labeling* of  $T$  is a bijection  $\ell : T \rightarrow \{1, 2, \dots, n\}$  such that  $\ell(v) \leq \ell(w)$  for all descendents  $w$  of  $v$ . Let  $f^T$  be the number of increasing labelings. The *hooklength*,  $h_v$ , of a vertex  $v$  is the number of descendents of  $v$ . The hook length formula for trees states that

$$f^T = \frac{n!}{\prod_{v \in T} h_v}.$$

There is a similar formula for the number of standard Young tableaux of given shape. Greene, Nijenhuis, and Wilf gave a beautiful probabilistic proof of the tableau formula where the hooklengths enter in a very natural way.

Recently, Han discovered a formula with the interesting property that hooklengths appear as exponents. Specifically, let  $\mathcal{B}(n)$  be the set of all  $n$ -vertex binary trees. Han proved algebraically that

$$\sum_{T \in \mathcal{B}(n)} \prod_{v \in T} \frac{1}{h_v 2^{h_v-1}} = \frac{1}{n!}.$$

We show how to give a simple probabilistic proof of this equation as well as various generalizations. We also pose some open questions raised by this work. (Received August 16, 2009)